

AD-A119 735

STATE UNIV OF NEW YORK AT BINGHAMTON DEPT OF MATHEMAT--ETC F/G 12/1
DETERMINING THE PROBABILITY OF AT LEAST ONE SUCCESS IN TRIALS C--ETC(U)
SEP 82 M YADIN, S.ZACKS

N00014-81-K-0407

UNCLASSIFIED

TR-3

NL

[]
[]
[]



END
DATE
FORMED
18-82
DTIC

AD A119735

(12)

DETERMINING THE PROBABILITY OF AT LEAST
ONE SUCCESS IN TRIALS CONDUCTED ON THE
LIGHTED PORTION OF A STAR SHAPED CURVE
SUBJECT TO A POISSON SHADOWING PROCESS

By

M. Yadin and S. Zacks

TECHNICAL REPORT NO. 3

September 1, 1982

Prepared under Contract
N00014-81-K-0407 (NT 042-276)
For the Office of Naval Research

S. Zacks, Principal Investigator

Reproduction is permitted for any use of the U.S. Government

DEPARTMENT OF MATHEMATICAL SCIENCES
STATE UNIVERSITY OF NEW YORK AT BINGHAMTON
BINGHAMTON, NEW YORK 13901

This document has been approved
for public release and sale; its
distribution is unlimited.

DTIC
SELECTE
SEP 29 1982
A

DTIC FILE COPY

82 09 077

DETERMINING THE PROBABILITY OF AT LEAST ONE SUCCESS IN TRIALS
CONDUCTED ON THE LIGHTED PORTION OF A STAR SHAPED CURVE SUBJECT
TO A POISSON SHADOWING PROCESS

By

M. Yadin and S. Zacks^{†)}

Technion, Israel Institute of Technology
and
State University of New York at Binghamton

Abstract

A star shaped curve, C , in the plane is subject to a Poisson shadowing process. According to this process, disks of random size appear at random locations in a region between a source of light, which is at the origin, and the curve C . These disks cast shadows on C . Trials are conducted along the lighted portion of C . Each trial requires a fixed length, ℓ , of C . The different trials are independent and have a fixed probability, p , of success. The number of trials conducted along C is a random variable, N , which depends on the random length of the lighted portion of C . The success probability is $P = 1 - E\{q^N\}$, where $q=1-p$. Lower and upper bounds for P are derived. A numerical example shows cases in which these bounds could be very close.

Key Words: Poisson Shadowing Process, Random Fields, Measure of Visibility, Moments of Visibility, Success Probabilities

^{†)} Research supported under contract N00014-81-K-0407 (NR 042-276) with the Office of Naval Research, at SUNY-Binghamton.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

1. Introduction

Consider a star-shaped curve in the plane, C , and a source of light at the origin, O . If there are no obstacles between the origin and the curve C , the whole curve is in the light or visible. Certain experiments (trials) can be conducted along the lighted portion of the curve. Each such trial requires a length ℓ of C , and the probability of its success is p , $0 < p < 1$. Let $L\{C\}$ be the total length of C . If C is completely visible, $N = [L\{C\}/\ell]$ trials can be conducted. ($[a]$ designates the integer part of a .) Assume that all these trials are independent, having the same success probability, p , (Bernoulli trials). Thus, the probability of at least one success, when C is completely lighted, is $S = 1 - q^N$, where $q = 1 - p$. In reality, C may not be completely visible, due to shadows cast on it by objects (disks, say), which are randomly dispersed in a region between O and C . The centers of the disks follow a Poisson process and their diameters are of random size. Thus, the total number of trials that can be performed along the visible portion of C is the random variable

$$N = \sum_{i=1}^K [X_i/\ell], \text{ where } K \text{ is the random number of connected subsets}$$

(disjoint segments) of C , which are visible, and X_i is the length of the i -th such subset. The probability of at least one success, under this random shadowing, is $P = 1 - Q$, where $Q = E\{q^N\}$. The present study develops a method for determining upper and lower bounds for Q . This method is based on the methodology developed by Yadin and Zacks [1], for determining the moments of the total visible portion of C ,

$$V\{C\} = \sum_{i=1}^k X_i. \text{ We show that a lower bound for } Q \text{ (an upper bound for}$$

P , respectively) is the value of the moment generating function (MGF) of $V\{C\}$, at the point $N = \ell nq/\ell$. The upper bound for Q (lower bound for P) can be obtained by considering $Q^* = E\{q^{N^*}\}$, where

$$N = \sum_{i=1}^k \left(\frac{X_i - \ell}{\ell} \right)_+, \text{ and } a_+^* = \max(a, 0). \text{ In section 2 we present the}$$

structure of the Poisson field of shadowing disks and review the main

results of [1], which lead to the MGF of $V\{C\}$. In Section 3 we present lower and upper bounds for Q , which are based on the MGF of $V\{C\}$ and

$$V^*\{C\} = \sum_{i=1}^K (X_i - \ell)_+.$$

Section 4 presents a numerical example for annular regions and a standard-uniform Poisson field of shadowing disks. It is shown, in the numerical example of Section 4, that the lower and upper bounds for Q developed in the present paper could be very close and effective. The present paper is very tightly linked with our previous paper [1], in which the general theory for the determination of the moments of visibility, $\mu_n = E\{V^n\{C\}\}$, is presented. As such it could be considered as an extension of [1] for an important class of applications.

2. The Poisson Field of Shadowing Obstacles, and the Moments of $V\{C\}$.

Consider a star-shaped curve, C , given by a function $r(s)$ on an interval $[s', s'']$, i.e.,

$$C = \{(\rho, \theta) ; \rho = r(\theta) , s' \leq \theta \leq s''\} . \quad (2.1)$$

We further assume that shadows on C are cast by disks, which are randomly distributed within a region, C_1 , bounded by the curves

$$U = \{(\rho, \theta) ; \rho = u(\theta) , s^* \leq \theta \leq s^{**}\} \text{ and}$$

$W = \{(\rho, \theta) ; \rho = w(\theta) , s^* \leq \theta \leq s^{**}\}$. Each disk is characterized by a point (ρ, θ, y) , where (ρ, θ) are the polar coordinates of its center and y is its diameter. It is assumed that the centers are uniformly distributed between U and W and the diameters of the disks are i.i.d. random variables having a c.d.f. $G(y)$ concentrated on $[a, b]$ (the standard case). Moreover, $u(\theta)$, $w(\theta)$ and b are such that both the origin, 0 , is uncovered and the curve C is not intersected by any one of the random disks. For the precise conditions see [1] . It is further assumed that the number of disks whose centers fall within a subset C of C_1 has a Poisson distribution, with mean A , when A is the area of C .

A point $P = (r(s), s)$ is said to be visible, if the ray \overline{OP} is not intersected by any shadowing disk.

The measure of visibility $V\{C\}$ is defined as

$$V\{C\} = \int_{s'}^{s''} I(s) \ell(s) ds , \quad (2.2)$$

where $I(s)=1$ if the point $(r(s), s)$ is in the light (visible), and $I(s)=0$ otherwise. $\ell(s)ds = [r^2(s) + (r'(s))^2]^{1/2}ds$ is the infinitesimal length of C at $(r(s), s)$. The moments of $V\{C\}$ were expressed in [1] in terms of the K-functions, $K_-(s, t)$ and $K_+(s, t)$. $\mu K_-(s, t)$ and $\mu K_+(s, t)$ are, respectively, the expected number of disks in C_1 , whose centers have orientation coordinates in $[s-t, s]$ ($[s, s+t]$, resp.), and which do not intersect the ray with orientation s . It is shown in [1] that these functions are given by

$$K_{-}(s, t) = \int_{s-t}^s \int_{u(\theta)}^{w(\theta)} G(y(\rho, s-\theta)) \rho d\rho d\theta$$

and

(2.3)

$$K_{+}(s, t) = \int_s^{s+t} \int_{u(\theta)}^{w(\theta)} G(y(\rho, \theta-s)) \rho d\rho d\theta,$$

where

$$y(\rho, \theta-s): y(\rho, s-\theta) = \begin{cases} 2\rho \sin|s-\theta| & , \text{ if } |s-\theta| < \pi/2 \\ 2\rho & , \text{ if } |s-\theta| \geq \pi/2 \end{cases} \quad (2.4)$$

is the maximal diameter of a disk centered at (ρ, θ) , which does not intersect the ray with orientation s .

It is shown in [1] that the n -th moment of $V\{C\}$ is

$$\mu_n = n! \int_{s' \leq s_1 \leq \dots \leq s_n \leq s''} \dots \int p(s_1, \dots, s_n) \prod_{i=1}^n \ell(s_i) ds_i, \quad (2.5)$$

where $p(s_1, \dots, s_n)$ is the probability that n points on C , with orientation coordinates s_1, \dots, s_n are simultaneously visible. It is further shown that

$$p(s_1, \dots, s_n) = \exp\{-V\{C_1\}\} \exp\{\mu K_{-}(s_1, s_1-s^*) +$$

(2.6)

$$\mu K_{+}(s_n, s_n^{**}-s_n) + \mu \sum_{i=1}^{n-1} \left(K_{+}(s_i, \frac{s_{i+1}-s_i}{2}) + K_{-}(s_{i+1}, \frac{s_{i+1}-s_i}{2}) \right) \},$$

in which

$$v\{C_1\} = \frac{\mu}{2} \int_{s^*}^{s^{**}} (w^2(s) - u^2(s)) ds \quad (2.7)$$

Furthermore, let

$$\psi_0(s) = \exp\{\mu K_-(s, s-s')\} \quad (2.8)$$

and define recursively, for $j \geq 1$

$$\psi_j(s) = \int_{s'}^s \ell(y) \psi_{j-1}(y) \exp\{\mu K_-(y, \frac{s-y}{2}) + \mu K_+(y, \frac{s-y}{2})\} dy \quad (2.9)$$

Then

$$\mu_n = n! \exp\{-v\{C_1\}\} \int_{s'}^{s''} \ell(s) \psi_{n-1}(s) \exp\{\mu K_+(s, s^{**}-s)\} ds \quad (2.10)$$

3. The MGF of $V\{C\}$ and The Bounds for Q

In Section 1 we introduced the random variables K, X_1, X_2, \dots, X_K , which are the number of lighted (visible) disjoint segments of C , and their length. Accordingly,

$$V\{C\} = \sum_{i=1}^K X_i. \text{ We also defined the random variable } N = \sum_{i=1}^K \left\lfloor \frac{X_i}{\ell} \right\rfloor.$$

Thus $N \leq V\{C\}/\ell$, with probability one. It follows, for every q , $0 < q < 1$, that

$$Q = E\{q^N\} \geq E\left\{q^{V\{C\}/\ell}\right\} = M_V(\ln q / \ell), \quad (3.1)$$

where $M_V(u)$ is the MGF of $V\{C\}$. Thus, $M_V(\ln q / \ell)$ is a lower bound for Q . Notice that, since $V(C) \leq L(C) < \infty$, all the moments of $V(C)$ are bounded by powers of $L(C)$. Hence, the MGF of $V(C)$ can be expressed as

$$M_V(u) = \sum_{i=0}^{\infty} \frac{u^i}{i!} \mu_i, \quad -\infty < u < \infty. \quad (3.2)$$

For the derivation of an upper bound for Q , we consider the random

variable $N^* = \sum_{i=1}^K \frac{X_i - \ell}{\ell} +$. Since $N^* < N$ with probability one,

$$Q^* = E\{q^{N^*}\} \geq Q. \quad (3.3)$$

In order to obtain Q^* we define a new visibility measure

$$V^*(C) = \sum_{i=1}^K (X_i - \ell)_+ \quad (3.4)$$

The moments of $V^*(C)$ can be obtained by the formulae presented in Section 2, in which the K -functions in (2.6) - (2.10) are modified in the following manner. Replace $K_-(s, t)$ and $K_+(s, t)$ by $K_-(s - \tau_1(s), (t - \tau_1(s))_+)$ and $K_+(s + \tau_1(s), (t - \tau_1(s))_+)$, respectively, where $\tau_1(s)$ should be determined so that

$$\int_{s - \tau_1(s)}^s \ell(s) ds = \ell/2 \quad (3.5)$$

and, similarly, $\tau_2(s)$ should satisfy the equation

$$\int_s^{s+\tau_2(s)} \ell(s) ds = \ell/2. \quad (3.6)$$

By definition, $K_{\pm}(s,0) = 0$ for all s .

More specifically, let μ_n^* ($n=1,2,\dots$) be the n -th moment of $V^*(C)$, which is given by

$$\mu_n^* = n! \int_S \dots \int p^*(s_1, \dots, s_n) \prod_{i=1}^n \ell(s_i) ds_i, \quad (3.7)$$

where $S = \{s' \leq s_1 \leq \dots \leq s_n \leq s''\}$.

The function $p^*(s_1, \dots, s_n)$ is the probability that the union of n segments of C , each one of length ℓ , centered around the points $(r(s_i), s_i)$, $i=1, \dots, n$, is completely visible. In other words, define the indicator function $I^*(s)$, which is equal to 1 if the segments of C of length ℓ , centered at $(r(s), s)$, is completely visible and is equal to 0 otherwise. Accordingly,

$$V^*(C) = \int_{s'}^{s''} I_{\ell}^*(s) \ell(s) ds. \quad (3.8)$$

As explained above, $p^*(s_1, \dots, s_n) = E\{\prod_{i=1}^n I^*(s_i)\}$. Following the theory developed in [1], $p^*(s_1, \dots, s_n)$ is given, as in (2.6), by

$$\begin{aligned} p^*(s_1, \dots, s_n) = & \exp\{-v\{C_1\}\} \exp\left\{\mu K_-\left(s_1 - \tau_1(s_1), (s_1 - \tau_1(s_1) - s)_+\right)\right. \\ & \left.+ \mu K_+\left(s_n + \tau_2(s_n), (s'' - s_n - \tau_2(s_n))_+\right) + \right. \\ & \left. \mu \sum_{i=1}^{n-1} \left(K_+\left(s_i + \tau_2(s_i), \left(\frac{s_{i+1} - s_i}{2} - \tau_2(s_i)\right)_+\right) + \right. \right. \\ & \left. \left. K_-\left(s_i - \tau_1(s_i), \left(\frac{s_{i+1} - s_i}{2} - \tau_1(s_i)\right)_+\right)\right)\right\}. \quad (3.9) \end{aligned}$$

Finally, if $M_{V^*}(u)$ denotes the MGF of $V^*(C)$, then $Q^* = M_{V^*}(\ln q/\ell)$, which is the upper bound for Q . In the following section we illustrate these bounds in a special case.

4. Lower And Upper Bounds For Q in A Special Case

In the present section we exhibit the method of determining lower and upper bounds for the failure probability Q in the following special case. The curve C is an arc on a circle of radius r , centered at the origin, limited by rays having orientations s' and s'' ,

$-\frac{\pi}{2} < s' \leq s'' \leq \frac{\pi}{2}$. The centers of the disks are distributed between U and W , where U and W are circles centered at the origin, with radii $0 < u < w < r$. Moreover, we assume that the centers of the disks are uniformly distributed within this annular region, and their random diameters, Y , are uniformly distributed between $[a, b]$ independently of their centers, where $0 < \frac{b}{2} \leq u < w \leq r - \frac{b}{2}$. This special case was previously studied in [1]. We have shown that in the present case, $K_-(s, t) = K_+(s, t) \equiv \hat{K}(t)$. Explicit formulae for this function can be found in [1]. Notice that in the present case of C being a circular arc, $\tau_1(s) = \tau_2(s) = \ell/2r$ for all s . Accordingly, in the determination of Q^* we replace $\hat{K}(t)$ by $\hat{K}(t - \frac{\ell}{2r})_+$. The computation of the moments μ_n and μ_n^* follows the procedure described in [1].

Let $\{\eta_n, n \geq 1\}$ be the normalized moments of $V\{C\}$, i.e.,

$\eta_n = E\{V^n(C)/r^n(s''-s')^n\}$. The sequence $\{\eta_n; n \geq 1\}$ is decreasing and, as shown in [1], $\lim_{n \rightarrow \infty} \eta_n = P_1$, which is the probability of complete

visibility of C . Furthermore, $M_V(\ln q/\ell) = \sum_{n=0}^{\infty} \frac{\theta^n}{n!} \eta_n$, where

$\theta = (\ln q) r(s''-s')/\ell$. One can approximate this MGF, to any degree of accuracy, in the following manner. Given an arbitrary $\epsilon, \epsilon > 0$, let m be a positive integer such that

$$\left(e^{|\theta|} - \sum_{i=0}^m \frac{|\theta|^i}{i!} \right) (\eta_m - P_1) \leq \epsilon. \quad (4.1)$$

Let

$$\hat{M}_V\left(\frac{\ln q}{\ell}\right) = \sum_{i=0}^{m-1} \frac{\theta^i}{i!} \eta_i + \eta_m \left(e^\tau - \sum_{i=0}^{m-1} \frac{\theta^i}{i!} \right). \quad (4.2)$$

Then, $\delta = \left| \hat{M}_V\left(\frac{\ln q}{\ell}\right) - M_V\left(\frac{\ln q}{\ell}\right) \right| \leq \epsilon$. Indeed,

$$\begin{aligned} \delta &= \left| \sum_{i=m+1}^{\infty} \frac{\theta^i}{i!} (\eta_m - \eta_i) \right| \leq \sum_{i=m+1}^{\infty} \frac{|\theta|^i}{i!} [\eta_m - \eta_i] \\ &\leq (\eta_m - p_1) \left(e^{|\theta|} - \sum_{i=0}^m \frac{|\theta|^i}{i!} \right) \end{aligned} \quad (4.3)$$

$M_{V*}\left(\frac{\ln q}{\ell}\right)$ can be approximated in the same manner. In Table 1 we provide numerical values of the lower and upper bounds for Q , corresponding to the following parameters of an annular region: $r=1.0$, $w=.6$, $u=.4$. The parameters of the distribution of Y are $a=.1$ and $b=.5$. These bounds are given for two values of μ , two values of ℓ , two values of $\Delta = s'' - s'$, and $q = .8$.

Table 1. Lower and Upper Bounds for Q , Circular Arc, C , And Annular Region of Disk Centers

	$\Delta = 120^\circ$		$\Delta = 60^\circ$	
	$\mu=1$	$\mu=5$	$\mu=1$	$\mu=5$
$\ell=.2$.126861	.258360	.353190	.502616
	.117760	.210320	.341425	.454504
$\ell=.4$.356616	.521052	.596354	.721755
	.337353	.439304	.580451	.661923

This table shows that in the present case the method developed here is very satisfactory.

REFERENCE

- [1] Yadin, M. and Zacks, S. (1982). Visibility Probabilities and Moments of Measures of Visibility On Star Shaped Curves In The Plane For Poisson Shadowing Processes.
Technical Report No. 2, ONR Contract N00014-81-K-0407 (NR 042-276) with SUNY-Binghamton.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 3	2. GOVT ACCESSION NO. AD-A119 735	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) DETERMINING THE PROBABILITY OF AT LEAST ONE SUCCESS IN TRIALS CONDUCTED ON THE LIGHTED PORTION OF A STAR SHAPED CURVE SUBJECT TO A POISSON SHADOWING PROCESS		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) M. Yadin and S. Zacks		6. PERFORMING ORG. REPORT NUMBER N00014-81-K-0407
9. PERFORMING ORGANIZATION NAME AND ADDRESS SUNY-Binghamton Binghamton, NY 13901		8. CONTRACT OR GRANT NUMBER(s) NR 042-276
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 --		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Office of Naval Research Arlington, VA 22217		12. REPORT DATE August 23, 1982
		13. NUMBER OF PAGES 12
		15. SECURITY CLASS. (of this report) non-classified
		16. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Poisson Shadowing Process, Random Fields, Measure of Visibility, Moments of Visibility, Success Probabilities		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A star shaped curve, C , in the plane is subject to a Poisson shadowing process. According to this process, disks of random size appear at random locations in a region between a source of light, which is at the origin, and the curve C . These disks cast shadows on C . Trials are conducted along the lighted portion of C . Each trial requires a fixed length, ℓ , of C . The different trials are independent and have a fixed probability, p , of success. The number of trials conducted along C is a random variable, N , which depends		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

S. N. 0102-LF-314-5601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. ABSTRACT (continued)

on the random length of the lighted portion of C . The success probability is $P = 1 - E\{q^N\}$, where $q = 1 - p$. Lower and upper bounds for P are derived. A numerical example shows cases in which these bounds could be very close.

ity is

A

1.

END

DATE
FILMED

11-8

DTIC